



# $g-2$ と 125GeV Higgs

[岩本 祥](#) [Sho IWAMOTO]

The University of Tokyo

9 Mar. 2012

@ 札幌ウインタースクール2012 (北大)

Based on

**M. Endo, K. Hamaguchi, S.I., N. Yokozaki. [1112.5653]**

Also See: Endo, Hamaguchi, SI, Yokozaki. [1108.3071] [1202.2751]  
Endo, Hamaguchi, SI, Nakayama, Yokozaki. [1112.6412]

# summary

To explain

$(g - 2)_\mu$  & 125 GeV Higgs  
simultaneously,

Extending the MSSM with  
vector-like quarks is

a very

attractive way.

# Standard Model

☹️ “hierarchy problem”

↓ SUSY around TeV

**MSSM** [Minimal Supersymmetric Standard Model]

😊 fermion/boson unification

😊 GUTs, dark matter(?)

😊 nicely explain **muon**  $g - 2$  anomaly

☹️ must be broken ... too many ~~SUSY~~ parameters



mSUGRA / GMSB frameworks



Physicists' triumph


Physicists' dream




# muon $g - 2$ anomaly

SM (HLMNT '11)

Expm (BNL '04)



Hagiwara, Liao, Martin, Nomura, Teubner [1105.3149]



- 😊 fermion/boson unification
- 😊 GUTs, dark matter(?)
- 😊 nicely explain muon  $g - 2$  anomaly
- 😞 must be broken ... too many ~~SUSY~~ parameters

↓  
mSUGRA / GMSB frameworks

# Standard Model

☹️ “hierarchy problem”

↓ SUSY around TeV


**MSSM** [Minimal Supersymmetric Standard Model]

😊 fermion/boson unification

😊 GUTs, dark matter(?)

😊 nicely explain  $\mu$ on  $g - 2$  anomaly

☹️ must be broken ... too many ~~SUSY~~ parameters



**mSUGRA / GMSB** frameworks

**However**

Now this “dream” is threatened by





Physicists' triumph



Physicists' dream



# The $\sim 125\text{ GeV}$ Higgs boson




# The $\sim 125\text{GeV}$ Higgs boson





# The $\sim 125\text{GeV}$ Higgs boson



$(g - 2)_\mu$  explanation in SUSY


# 125 GeV in MSSM

$$m_h^2 \lesssim m_Z^2 + \frac{3g_W^2 m_t^4}{8\pi^2 m_W^2} \left[ \ln \frac{M_S^2}{m_t^2} + \alpha^2 \left( 1 - \frac{\alpha^2}{12} \right) \right]$$

(1-loop level)

where  $M_S^2 := \frac{M_{t_1}^2 + M_{t_2}^2}{2}$ ,  $\alpha := \frac{A_t - \mu \cot \beta}{M_S}$ .

- heavy  $\tilde{t}$
- large  $(A_t - \mu \cot \beta)$   
(roughly  $\approx -\sqrt{6}m_{\tilde{t}}$ )




# $(g - 2)_\mu$ in MSSM

$$\Delta(\tilde{\chi}^\pm, \tilde{\nu}) \approx \frac{\alpha_w m_\mu^2}{m_{\text{soft}}^2} \text{sgn}(\mu M_2) \tan \beta,$$

$$\Delta(\tilde{\chi}^0, \tilde{\mu}) \approx \frac{\alpha_Y m_\mu^2}{m_{\text{soft}}^2} \text{sgn}(\mu M_1) \tan \beta + \dots$$

- light  $(\tilde{\nu}_\mu, \tilde{\chi}^\pm)$  or  $(\tilde{\mu}, \tilde{\chi}^0)$
- large  $\tan \beta$



[10]

# 125 GeV in MSSM

$$m_h^2 \lesssim m_Z^2 + \frac{3g_W^2 m_t^4}{8\pi^2 m_W^2} \left[ \ln \frac{M_S^2}{m_t^2} + \alpha^2 \left( 1 - \frac{\alpha^2}{12} \right) \right]$$

(1-loop level)

where  $M_S^2 := \frac{M_{t_1}^2 + M_{t_2}^2}{2}$ ,  $\alpha := \frac{A_t - \mu \cot \beta}{M_S}$ .

- heavy  $\tilde{t}$

dilemma (GUT)



- large  $(A_t - \mu \cot \beta)$   
(roughly  $\approx -\sqrt{6}m_{\tilde{t}}$ )

# $(g - 2)_\mu$ in MSSM

$$\Delta(\tilde{\chi}^\pm, \tilde{\nu}) \approx \frac{\alpha_w m_\mu^2}{m_{\text{soft}}^2} \text{sgn}(\mu M_2) \tan \beta,$$

$$\Delta(\tilde{\chi}^0, \tilde{\mu}) \approx \frac{\alpha_Y m_\mu^2}{m_{\text{soft}}^2} \text{sgn}(\mu M_1) \tan \beta + \dots$$

- light  $(\tilde{\nu}_\mu, \tilde{\chi}^\pm)$  or  $(\tilde{\mu}, \tilde{\chi}^0)$

- large  $\tan \beta$

GMSB ... small  $A$ -terms  $\implies$  impossible!

## mSUGRA

- $b \rightarrow s\gamma$  forbids a huge  $A$ -term [1112.6412]
- large  $\mu \tan \beta$  + small  $m_{\tilde{\tau}}$   $\implies$  instable vacuum etc...  
Hisano, Sugiyama [1011.0260]

$\implies$  Possible with parameter splitting & tuning.

e.g. "Non-Universal Gaugino Model" or to split  $M_0$  for 1,2 / 3 gen.



[11]

# For 125 GeV & $g-2$ , we must...

- ◎ **tune** the parameter in ~~SUSY~~ models
- ◎ **ignore**  $(g - 2)_\mu$  anomaly.
  - “It is just from hadronical uncertainty, theorists’ fault!!”
- ◎ **wish** a lighter Higgs.
- ◎ **extend** the MSSM.
  - NMSSM
  - add  $5 + \bar{5}$
  - add  $10 + \overline{10}$
  - add a new gauge symmetry.

# For 125 GeV & $g-2$ , we must...

- ◎ **tune** the parameter in ~~SUSY~~ models
- ◎ **ignore**  $(g-2)_\mu$  anomaly.
  - “It is just from hadronical uncertainty, theorists’ fault!”
- ◎ **wish** a lighter Higgs.
- ◎ **extend** the MSSM.
  - ~~NMSSM~~  $g-2 \Rightarrow$  large  $\tan\beta \Rightarrow$  NMSSM not contribute.
  - ~~add  $5 + \bar{5}$~~  is still inadequate. Martin [0910.2732]
  - **add  $10 + \bar{10}$**  **Today’s topic.** [1112.5653]
  - add a new gauge symmetry. ← See: Endo, Hamaguchi, SI, Nakayama, Yokozaki [1112.6412]

## 2. The Extension we propose

# Extension w. Vector-like Matters

$$\text{MSSM} + (\mathbf{10} + \overline{\mathbf{10}}), \text{ i.e. } \begin{cases} \mathbf{10} = (Q', U', E') \\ \overline{\mathbf{10}} = (\bar{Q}', \bar{U}', \bar{E}') \end{cases}$$

$$W_{\text{add}} = \overset{m_h \uparrow}{Y' Q' H_u U'} + Y'' \bar{Q}' H_d \bar{U}' \\ + M_V Q' \bar{Q}' + M_V U' \bar{U}' + M_V E' \bar{E}'$$

## IDEA

MSSM: top (s)quark lifts up higgs. Okay, then...

Add another top quark!


⇒ Gauge anomaly...

⇒ Add as vector-like!

# Extension w. Vector-like Matters

$$\text{MSSM} + (\mathbf{10} + \overline{\mathbf{10}}), \text{ i.e. } \begin{cases} \mathbf{10} = (Q', U', E') \\ \overline{\mathbf{10}} = (\bar{Q}', \bar{U}', \bar{E}') \end{cases}$$

- No gauge anomaly.
- Gauge couplings unification.



Martin [0910.2732]

$$W_{\text{add}} = \overset{m_h \uparrow}{Y'} Q' H_u U' + \overset{m_h \downarrow}{Y''} \bar{Q}' H_d \bar{U}' + M_V Q' \bar{Q}' + M_V U' \bar{U}' + M_V E' \bar{E}'$$

$m_h \downarrow \rightarrow \text{we assume } Y'' \ll 1.$

$$W_{\text{mix}} = \epsilon_i Q_i H_u U' + \epsilon'_i Q' H_u \bar{U}_i + \epsilon''_i Q' H_d \bar{D}_i$$

## Mixing between SM- & vector-like quark

- Too large  $\rightarrow$  flavor problem?
- No mixing  $\rightarrow$  stable colored particle.

$\Rightarrow$  *assumed very small.*

params:  $(\Lambda, M_{\text{mess}}, \tan \beta, N_{\text{mess}}, \text{sgn } \mu, Y', M_V)$

(GMSB framework)

- $N_{\text{mess}} = 1$  to keep perturbative up to  $M_{\text{GUT}}$ .
- $\text{sgn } \mu = +$  to explain  $g - 2$ .
- $Y' = 1.05$  : infrared fixed point  $\Rightarrow$  nice for 125 GeV (also  $A_t$  and  $A'$  go to IR fixed point.)



# RESULT

in this talk

with  $\left\{ \begin{array}{l} \text{GMSB framework} \\ \text{mSUGRA framework} \end{array} \right.$



params:  $(\Lambda, M_{\text{mess}}, \tan \beta, N_{\text{mess}}, \text{sgn } \mu; Y', M_V)$

$\parallel$   
1

$\parallel$   
+

$\parallel$   
1.05

$$M_{\text{mess}} = 10^6 \text{ GeV}$$



simultaneous realization:


$$(M_V, m_{\tilde{g}}, \tan \beta) \approx$$

$$(600 \text{ GeV}, 900 \text{ GeV}, 20)$$

$$- (1000 \text{ GeV}, 1200 \text{ GeV}, 30)$$

$g - 2$   $1\sigma$   $2\sigma$   
 $124 \text{ GeV} < m_h < 126 \text{ GeV}$


params:  $(\Lambda, M_{\text{mess}}, \tan\beta, N_{\text{mess}}, \text{sgn}\mu; Y', M_V)$   
 $\parallel$   $\parallel$   $\parallel$   
 $1$   $+$   $1.05$



- $(g - 2)_\mu$  expm. tells us ( $2\sigma$ -level),  
 $M_V \lesssim 1.5 \text{ TeV}, m_{\tilde{g}} \lesssim 1.6 \text{ TeV}$
- If we take  $(g - 2)_\mu$  seriously ( $1\sigma$ -level),  
 $M_V \lesssim 1.0 \text{ TeV}, m_{\tilde{g}} \lesssim 1.2 \text{ TeV}$

# 3. LHC Phenomenology

# SUSY search



	prompt decay ( $M_{\text{mess}} \lesssim 10^5 \text{ GeV}$ )	long-lived ( $M_{\text{mess}} \gtrsim 10^6 \text{ GeV}$ )
$\tilde{\chi}^0$	curr. ( $1\text{fb}^{-1}$ ) $m_{\tilde{g}} \gtrsim 1.2 \text{ TeV}$ $2\gamma + \cancel{E}$ : ATLAS[1111.4116]	curr. ( $1\text{fb}^{-1}$ ) $m_{\tilde{g}} \gtrsim 700 \text{ GeV}$ $4j + \cancel{E}$ : ATLAS1109.6572, CMS SUS11-008
$\tilde{\tau}$	will easily be covered by multi-lepton search	already excluded (CMS hvy-stbl chrgd prtcl; Seminar Jan. '12)

# Vector-like Quark Search

**direct proof!**

- ◉ New “vector-like” quark  $(t'_1, b', t'_2)$

$$10 = (Q', U', E')$$

$$\bar{10} = (\bar{Q}', \bar{U}', \bar{E}')$$

## Mass

$$m_{t'} \sim M_V \pm (174 \text{ GeV}/2),$$

$$m_{b'} = M_V$$

$$W_{\text{add}} = Y' Q' H_u U' + Y'' \bar{Q}' H_d \bar{U}'$$

$$+ M_V Q' \bar{Q}' + M_V U' \bar{U}' + M_V E' \bar{E}'$$


$$W_{\text{mix}} = \epsilon_i Q_i H_u U' + \epsilon'_i Q' H_u \bar{U}_i + \epsilon''_i Q' H_d \bar{D}_i$$

depending on mixing  
btw. vec-like/SM quark.

## Production

$$pp \rightarrow t'_1 \bar{t}'_1 \text{ etc. (pair production)}$$

## Decay



# Vector-like Quark Search

- ◉ New “vector-like” quark ( $t'_1, b', t'_2$ )

## Current bound

$$pp \rightarrow t'_1 \bar{t}'_1; \quad t'_1 \begin{cases} \rightarrow qW \\ \rightarrow qZ \\ \rightarrow qh \quad (\rightarrow q b \bar{b}) \end{cases}$$

if it decays exclusively as

$$t'_1 \rightarrow bW \quad :: m_{t'_1} > 552 \text{ GeV} \quad \text{CMS } 4.7\text{fb}^{-1} \text{ [EXO-11-050]}$$

$$t'_1 \rightarrow q_d W \quad :: m_{t'_1} > 340 \text{ GeV} \quad \text{CDF } 5.6\text{fb}^{-1} \text{ [1107.3875]}$$

$$t'_1 \rightarrow tZ \quad :: m_{t'_1} > 475 \text{ GeV} \quad \text{CMS } 1.14\text{fb}^{-1} \text{ [1109.4985]}$$

$$t'_1 \rightarrow q_u Z \quad :: \text{No bound yet}$$

$$t'_1 \rightarrow th \quad :: \text{No bound yet}$$

$$t'_1 \rightarrow q_u h \quad :: \text{No bound yet}$$


} No general bound on  $t'_1$  yet because of these possibility.

$$\geq 4 \text{ } b\text{-quarks} \quad (h \rightarrow b \bar{b})$$


Interesting channel after Higgs discovery.

# Vector-like Quark Search

At LHC **8**TeV,  
production **doubles!**



More severe bounds,  
or....?



$t'_1 \rightarrow th$  :: **No bound yet**

$t'_1 \rightarrow q_u h$  :: **No bound yet**

No general bound on  $t'_1$  yet  
because of these possibility.

$\geq 4$   $b$ -quarks ( $h \rightarrow b\bar{b}$ )


Interesting channel after Higgs discovery.




# 4. Conclusion

# Conclusion

125 GeV higgs? +  $(g - 2)_\mu$



MSSM +  $\mathbf{10} + \overline{\mathbf{10}}$  : vector-like quarks



Our ~~delusion~~ dream will be smashed/proved by

- SUSY search ( $\tilde{\chi}_1^0$ -(N)LSP /  $\tilde{\tau}$ -(N)LSP)
- 4th gen. quark search
  - $t' \rightarrow qW$
  - $t' \rightarrow qZ$
  - $t' \rightarrow qh (\rightarrow q_u b \bar{b})$

at the LHC