

# MSSM + $(g-2)_\mu$ + LHC

Sho IWAMOTO (岩本 祥)

21 Aug. 2018

Beyond Standard Model: Where do we go from here? @ GGI

Based on

- Endo, Hamaguchi, Iwamoto, Yanagi [[1704.05287](#)]
- Endo, Hamaguchi, Iwamoto, Yoshinaga [[1303.4256](#)]

and a few ongoing projects.

(cf. [https://github.com/misho104/notes/blob/master/MSSM/journal\\_20180313.pdf](https://github.com/misho104/notes/blob/master/MSSM/journal_20180313.pdf))

**1.  $(g-2)_\mu$**

**2. LHC**

LHC:  $\sim 150 \text{ fb}^{-1}$  (2018)  
 $+ 150 \text{ fb}^{-1}$  (2021-23)

$\begin{matrix} g-2 \\ 2021 \\ \sim \end{matrix}$

SUSY (MSSM): Solution to

- Naturalness if

$$\begin{aligned} H &\lesssim 200 \text{ GeV} \\ \tilde{t} &\lesssim 600 \text{ GeV} \\ \tilde{g} &\lesssim 900 \text{ GeV} \end{aligned} \quad \left. \right) \text{ for } 20\% \text{ tuning} \quad [1110.6926]$$

mostly excluded (except for degenerated region)

- DM

- Gauge coupling unification

- muon  $g-2$  anomaly

$$a_{\mu}^{\text{SM}} = \frac{g_{\mu}^{\text{SM}} - 2}{2} = (11659182.04 \pm 3.56) \times 10^{-10} \quad [1802.02995]$$

BNL 2004:  $209.1 \pm 6.3$


$\rightarrow 3.7 \sigma$

Fermilab (ongoing):  $?? \pm \sim 6$  data collected (release 2019?)

$?? \pm 1.6$  (2020)  $\rightarrow 7 \sigma ??$

$(g-2)_\mu$  from SUSY

$$\text{Re} \left[ \frac{\mu}{M} \frac{1}{(1 - \frac{\mu}{M})} \right]$$

three SUSY particles  $\lesssim 1 \text{ TeV}$ 

$$\textcircled{1} \propto g_2^2 \times \frac{m_\mu^2 M_2 \mu \tan\beta}{m_{\text{Loop}}^4} \propto \text{sign}(\mu) \quad m_{\text{Loop}} = \begin{cases} m_{\tilde{\mu}_L}, m_{\tilde{\mu}_R}, m_{\tilde{W}} \\ S_S \\ \mu M_2 \end{cases}$$

$$\textcircled{2} = -\frac{1}{2} \times \textcircled{1} \quad \therefore \textcircled{1} + \textcircled{2} > 0 \text{ for } \mu > 0$$

$$\Rightarrow \mu > 0, \tan\beta = O(10)$$

$$\tilde{W}, \tilde{H}, \tilde{\mu} \lesssim 500-1 \text{ TeV}_{\text{GU}}$$

Wino  
Scenario

$$\textcircled{3} \quad \alpha - g_Y^2 \times \frac{m^2 M_1 M \tan \beta}{m_{\text{loop}}^4} \Rightarrow \mu < 0, \tan \beta = O(10),$$

$\tilde{B}, \tilde{H}, \tilde{M}_R \lesssim 400 \text{ GeV}$

$$\textcircled{4} \quad \alpha + g_Y^2 \times \frac{M_1 M \tan \beta}{m_{\text{loop}}^4} \quad \text{"negative } \mu \text{" because } U(1) \text{ Scenario}$$

$$\textcircled{5} \quad \alpha + g_Y^2 \times \frac{M_1 M \tan \beta}{m_{\text{loop}}^4} \quad \text{but } \mu \notin m_{\text{loop}} = \{M_L, M_R, M_1\}$$

$$\Rightarrow \alpha \mu \tan \beta \quad \mu = +O(\text{TeV})$$

$\tilde{B}, \tilde{M}_L, \tilde{M}_R \lesssim 1 \text{ TeV}$

pure-bino  
( $\mu \tan \beta$ -enhancement)

LHC?


	500GeV	1TeV
$p p \rightarrow \tilde{W}^+ \tilde{W}^0$	33 fb	1.0
$\tilde{W}^+ \tilde{W}^-$	22	0.62
$\tilde{H}^+ \tilde{H}^0$	11	0.3
$\tilde{H}^+ \tilde{H}^-$	6.2	0.18
$\tilde{l}_L \tilde{l}_L^*$	0.47	
$\tilde{l}_R \tilde{l}_R^*$	0.18	

pure-bino  $\rightarrow p p \rightarrow \tilde{l}_{LR} \tilde{l}_{LR}^* \rightarrow 2l + E_T$

BHR "negative  $\mu$ "  $\rightarrow p p \rightarrow \tilde{H}^+ \tilde{H}^0 \rightarrow 3T + E_T, 3l + E_T$

Wino  $\rightarrow \tilde{W}^+ \tilde{W}^- \rightarrow 3l + E_T$



## SUSY contribution to muon $g-2$ : (1) "Chargino" contributions






$$[C] \quad \frac{g_2^2 m_\mu^2}{8\pi^2} \frac{M_2 \mu \tan \beta}{m_{\tilde{\nu}_\mu}^4} \cdot F_a \left( \frac{M_2}{m_{\tilde{\nu}_\mu}}, \frac{\mu}{m_{\tilde{\nu}_\mu}} \right)$$

$$[C'] \quad - \frac{g_2^2 m_\mu^2}{16\pi^2} \frac{M_2 \mu \tan \beta}{m_{\tilde{\mu}_L}^4} \cdot F_b \left( \frac{M_2}{m_{\tilde{\mu}_L}}, \frac{\mu}{m_{\tilde{\mu}_L}} \right)$$

**C+C'-contribution** [tree-level; slep=sneu]





## SUSY contribution to muon $g-2$ : gauge basis



# SUSY contribution to muon $g-2$ : (3) pure-Bino contribution


Endo, Hamaguchi, Kitahara, Yoshinaga [1309.3065]





**1.  $(g-2)_\mu$**

**2. LHC**

July 2018









$pp \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_1^\pm$


Moriond 2017






$pp \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ 

Moriond 2017




$pp \rightarrow \tilde{\chi}^0 \tilde{\chi}^+ (\widetilde{W}^0 \widetilde{W}^+ \text{ or } \widetilde{H}^0 \widetilde{H}^+)$ ; then?



$$\tilde{\chi}_2^0 \tilde{\chi}_1^+ \rightarrow ZW/hW + \text{mET}$$

$$(\rightarrow 3\ell + \text{mET})$$




but  $Z$ -like leptons




$$\tilde{\chi}_2^0 \tilde{\chi}_1^+ \rightarrow 3\ell + \text{mET}$$

Z-unlike


$pp \rightarrow \tilde{\chi}^0 \tilde{\chi}^+ (\widetilde{W}^0 \widetilde{W}^+ \text{ or } \widetilde{H}^0 \widetilde{H}^+)$ ; then?




$$x_\ell \sim 0.5$$




$$x_\ell \sim 1$$




$$x_\ell \sim 0$$





pMSSM w.  
 $\tilde{q}, \tilde{g}$ -decoupled.  
 $\tilde{\ell}_R, \tilde{\tau}_L, \tilde{\tau}_R$  also  
decoupled.

- $\tan \beta = 40$
- $M_1 = M_2/2$
- $\mu = M_2$



6 years ago!

- pMSSM w.  
 $\tilde{q}, \tilde{g}$ -decoupled.
- $\tilde{\ell}_R, \tilde{\tau}_L, \tilde{\tau}_R$  also  
decoupled.

- $\tan \beta = 40$
- $M_1 = M_2/2$
- $\mu = M_2$

# Backup

## Muon $g-2$ SM expectation : $3-4\sigma$ discrepancy!

$$a_\mu^{\text{SM}} \simeq \begin{array}{c} \text{(5-loop)} \\ \text{QED} \end{array} + \begin{array}{c} \text{(2+loop)} \\ \text{W,Z,H} \end{array} + \begin{array}{c} \text{(dispersion rel.)} \\ \text{had. vac. polarization} \end{array} + \begin{array}{c} \text{(low-energy EFT)} \\ \text{had. light-by-light} \end{array}$$

$$a_\mu(\text{QED}) = (11\,658\,471.886 \pm 0.003) \times 10^{-10},$$

$$a_\mu(\text{EW}) = ( -15.36 \pm 0.11 ) \times 10^{-10},$$

$$a_\mu(\text{HVP-LO}) = ( \quad 689.46 \pm 3.25 \quad ) \times 10^{-10},$$

$$a_\mu(\text{HVP-HO}) = (-8.70 \pm 0.07) \times 10^{-10},$$

$$a_\mu(\text{HLbL}) = ( -10.34 \pm 2.88 ) \times 10^{-10}.$$

$$a_\mu(\text{NP})? \dots 10 \times 10^{-10} \approx \frac{a_{\text{em}}}{4\pi} \left( \frac{m_\mu}{200 \text{ GeV}} \right)^2$$

$$a_{\mu}^{\text{SM}} = (11\,659\,178.3 \pm 4.3) \times 10^{-10}$$

$$a_{\mu}^{\text{SM}} = (11\,659\,209.2 \pm 6.3) \times 10^{-10}$$

(BNL '04+CODATA '14)

→ ±1.6 @ Fermilab in 1–2 year!

See also:

QED: Laporta [1704.06996], Marquard et al. [1708.07138].

HVP-LO: Keshavarzi, Nomura, Teubner [1802.02995]


HVP-HO: Kurz, Liu, Marquard, Steinhauser [1403.6400]

HIL No. KARZ, Elia, Marquard, Steinmäder [

Hebele, Segenhofer, Nyffeler [0902.5500],  
Colangelo, Hoferichter, Nyffeler, Passera, Stoffer [1403.7514]

## ■ Muon $g-2$ anomaly: What is the origin?

- Just a statistical fluctuation. *we assume it is "actual".*
- Just an issue in the experiment.
- $O(100)$  GeV particles with  $O(0.1)$  couplings
  - KK gravitino, MSSM
- keV–MeV particles with tiny couplings.
  - dark photon (extra U(1) gauge boson)



$$10 \times 10^{-10} \approx \frac{a_{\text{em}}}{4\pi} \left( \frac{m_\mu}{m_{\text{new}}} \right)^2$$

$\sim 200 \text{ GeV}$

$$10 \times 10^{-10} \approx \frac{(\epsilon^2/4\pi)}{4\pi} \left( \frac{m_\mu}{m_{\text{new}}} \right)^2$$

$\text{keV-MeV}$

$e^+ e^- \rightarrow \gamma A'$ ,  $A' \rightarrow \text{invisible}$   
BaBar [1702.03327]

## ■ Muon $g-2$ anomaly: What is the origin?

- Just a statistical fluctuation. we assume it is "actual".
- Just an issue in experiment.
- $O(100)$  GeV particles with  $O(0.1)$  couplings
  - MSSM
- keV–MeV particles with tiny couplings.
  - dark photon (extra U(1) gauge boson)
  - extra  $L_\mu - L_\tau$  gauge boson




$$10 \times 10^{-10} \approx \frac{a_{\text{em}}}{4\pi} \left( \frac{m_\mu}{m_{\text{new}}} \right)^2$$

$\sim 200$  GeV

$$10 \times 10^{-10} \approx \frac{(\varepsilon^2/4\pi)}{4\pi} \left( \frac{m_\mu}{m_{\text{new}}} \right)^2$$

keV–MeV


Gninenko, Krasnikov [[ph/0102222](#)],  
Baek, Deshpande, He, Ko [[ph/0104141](#)]



$$L_{Z'} = e_\mu Z'_\nu [\bar{\mu}\gamma^\nu\mu - \bar{\tau}\gamma^\nu\tau + \bar{\nu}_\mu\gamma^\nu\nu_\mu - \bar{\nu}_\tau\gamma^\nu\nu_\tau]$$

Gninenko, Krasnikov [[1801.10448](#)]

# SUSY contribution to muon $g-2$ : gauge basis



[C]	$\frac{g_2^2 m_\mu^2}{8\pi^2} \frac{M_2 \mu \tan \beta}{m_{\tilde{\nu}_\mu}^4} \cdot F_a \left( \frac{M_2}{m_{\tilde{\nu}_\mu}}, \frac{\mu}{m_{\tilde{\nu}_\mu}} \right)$
[C']	$-\frac{g_2^2 m_\mu^2}{16\pi^2} \frac{M_2 \mu \tan \beta}{m_{\tilde{\mu}_L}^4} \cdot F_b \left( \frac{M_2}{m_{\tilde{\mu}_L}}, \frac{\mu}{m_{\tilde{\mu}_L}} \right)$
[B]	$\frac{g_Y^2 m_\mu^2}{8\pi^2} \frac{\mu \tan \beta}{M_1^3} \cdot F_b \left( \frac{m_{\tilde{\mu}_L}}{M_1}, \frac{m_{\tilde{\mu}_R}}{M_1} \right)$
[BHR]	$-\frac{g_Y^2 m_\mu^2}{8\pi^2} \frac{M_1 \mu \tan \beta}{m_{\tilde{\mu}_R}^4} \cdot F_b \left( \frac{M_1}{m_{\tilde{\mu}_R}}, \frac{\mu}{m_{\tilde{\mu}_R}} \right)$
[BHL]	$\frac{g_Y^2 m_\mu^2}{16\pi^2} \frac{M_1 \mu \tan \beta}{m_{\tilde{\mu}_L}^4} \cdot F_b \left( \frac{M_1}{m_{\tilde{\mu}_L}}, \frac{\mu}{m_{\tilde{\mu}_L}} \right)$





$F_a, F_b$  are loop functions and positive.

$$\left. \begin{aligned} F_a(x, y) &= \frac{1}{2} \frac{C_1(x^2) - C_1(y^2)}{x^2 - y^2}, & F_b(x, y) &= -\frac{1}{2} \frac{N_2(x^2) - N_2(y^2)}{x^2 - y^2}; \\ C_1(x) &= \frac{3 - 4x + x^2 + 2 \log x}{(1-x)^3}, & N_2(x) &= \frac{1 - x^2 + 2x \log x}{(1-x)^3}. \end{aligned} \right\}$$

■ Wino  $\gg$  TeV & Higgsino  $<$  TeV  $\rightarrow$  BHL or BHR scenario.

$(\mu > 0)$        $(\mu < 0)$

- $\propto g_Y^2 \rightarrow$  relevant particles  $\lesssim 500$  GeV
- LHC:  $pp \rightarrow \tilde{H}^+ \tilde{H}^0, \tilde{H}^+ \tilde{H}^-$  "not much, but enough"
- DM: ~~Bino-Higgsino mixing~~ excl. by XENON1T bino-slepton co-annihilation, resonance. → future works



■ Bino-slepton (stau) co-annihilation  $\rightarrow m_{\tilde{\nu}_\tau}$  (or  $m_{\tilde{\tau}_R}$ )  $\simeq m_{\tilde{B}}$ .



■ We assumed:


- slepton universality,
- DM density is realized at each point in the plots.

$$\rightarrow m_{\tilde{B}} \simeq (m_{\tilde{\tau}_R} \text{ or } m_{\tilde{\nu}_\tau}) \lesssim m_{\tilde{\mu}} < m_{\tilde{H}} \quad (\sim M_1)$$

■ HL-LHC?

- $p p \rightarrow \tilde{H}^+ \tilde{H}^0, \tilde{H}^+ \tilde{H}^-$
- $\tilde{H}^0 \rightarrow \tau \tilde{\tau}, \tilde{H}^+ \rightarrow \tau \tilde{\nu}_\tau$   
due to  $\tan\beta$  + mass spectrum
- multi-tau signature







$\rightarrow m_{\tilde{B}} \simeq (m_{\tilde{\tau}_R} \text{ or } m_{\tilde{\nu}_\tau}) \lesssim m_{\tilde{\mu}} < m_{\tilde{H}} \text{ } (\sim M_1)$

## ■ HL-LHC?

- $pp \rightarrow \tilde{H}^+ \tilde{H}^0, \tilde{H}^+ \tilde{H}^-$
- $\tilde{H}^0 \rightarrow \tau \tilde{\tau}, \tilde{H}^+ \rightarrow \tau \tilde{\nu}_\tau$   
due to  $\tan\beta +$  mass spectrum
- multi-tau signature  
**"2 $\tau$  (+ soft) + missing"**



$(\mu > 0)$  $(\mu < 0)$ 

■ Wino  $\gg$  TeV & Higgsino  $<$  TeV  $\rightarrow$  BHL or BHR scenario.

- DM: Bino–stau co-annihilation  $\rightarrow m_{\tilde{B}} \simeq (m_{\tilde{\tau}_R} \text{ or } m_{\tilde{\nu}_\tau}) \lesssim m_{\tilde{\mu}} < m_{\tilde{H}}$
- DM has small Higgsino component  $\rightarrow$  LUX/XENON1T constraint.
- LHC:  $pp \rightarrow \tilde{H}^+ \tilde{H}^0, \tilde{H}^+ \tilde{H}^-; \tilde{H} \rightarrow \tau + \dots$  "2 $\tau$ +missing" signature

